



# **NEUTRA-STAT® TB**

## **READY TO USE DISINFECTANT CLEANER**

Effective against MRSA, VRE and Influenza A: H1N1\*\*\*

**HEALTHIER SURROUNDINGS:** Reduces cross-contamination.

**SAVES LABOR:** Cleans and disinfects in one step.

**SAFE TO USE:** No concentrate to handle.

**EASY TO USE:** No measuring and mixing.

**NEUTRAL pH:** Safe on all surfaces.

Neutra-Stat TB® is a ready-to-use "one-step" hospital grade disinfectant cleaner. It is tuberculocidal, bactericidal, virucidal and fungicidal.

Cross-contamination is a major housekeeping concern not only in hospitals, but in schools, institutions and industry. Neutra-Stat TB has been designed to aid in the reduction of cross-contamination in these areas. Neutra-Stat TB is effective against a broad spectrum of viral and bacterial pathogens including HIV-1 and Pseudomas aeruginosa. It is effective in the presence of 5% serum contamination and 400ppm water hardness. The end result is healthier surroundings when Neutra-Stat TB is used.

### **Saves labor:**

Neutra-Stat TB is not only a disinfectant, but a good cleaner as well. Using one product instead of two saves labor and reduces inventory. In addition Neutra-Stat TB will control and inhibit mold and mildew growth. As an added value, it may be used as a non-acid toilet bowl cleaner and disinfectant.

### **Deodorizes with a pleasant fragrance:**

Neutra-Stat TB contains a pleasant floral fragrance. The fragrance was chosen specifically to be non-offensive

to anyone's sense of smell. Neutra-Stat TB kills odor-causing bacteria where their growth causes malodors which enhances the image of a facility.

### **Easy to use:**

Neutra-Stat TB requires no mixing making it easy to use and safe to handle. Workers never come into contact with corrosive or irritating concentrates. Neutra-Stat TB is safe to use on any hard surface making it very versatile.

### **Directions:**

Apply undiluted to the surface using a sprayer, cloth or mop so as to thoroughly wet the surface. For complete disinfection the surface must remain wet for ten minutes.

### **Packaging:**

Neutra-Stat TB is packaged in quart containers, 12 quarts per case. For convenience, two sprayers are included in each case. All packaging is recyclable.

### **Product Data:**

Appearance: Colorless liquid

Odor: Floral

pH: 10.5 - 12.5

Stability: Stable from 32° F. to 105° F. for up to two years.

**Tuberculocidal Test Results:** Effective against Mycobacterium bovis according to the A.O.A.C. Confirmative In Vitro Test for Determining Tuberculocidal Activity in the presence of 5% organic serum with a 5 minute wet time.

**Bactericidal Test Results:** Effective against the following pathogenic bacteria according to the A.O.A.C. Germicidal Spray Products as Disinfectants test method, modified in the presence of 5% organic serum. Ten minute contact time.

Methicillin Resistant Staphylococcus aureus (MRSA)  
Vancomycin Resistant Enterococcus Faecalis (VRE)  
Vancomycin Intermediate Resistant Staphylococcus Aureus (VISA)  
Methicillin Resistant Staphylococcus epidermidis (MRSE)

Pseudomonas aeruginosa      Staphylococcus aureus  
E. coli and E. coli O157:H7      Salmonella typhi  
Salmonella Choleraesuis      Listeria monocytogenes  
Yersinia enterocolitica      Enterococcus Faecium  
Corynebacterium ammoniagenes  
Streptococcus pyogenes (Necrotizing Fasciitis-Group A)

**Fungicidal Test Results:** Fungicidal against Trichophyton interdigitale according to the A.O.A.C. Germicidal Spray Products as Disinfectants test method, modified in the presence of 5% organic serum. Ten minute contact time.

**Mildewstat Test Results:** Mildewstatic for Aspergillus niger according to the E.P.A. Hard Surface Mildew Fungistatic Test.

**Virucidal Test Results:** Virucidal against Hepatitis A, Canine Parvovirus and Poliovirus Type 1 with a wet time of 10 minutes; Hepatitis B, Hepatitis C, and Bovine Viral Diarrhea Virus with a wet time of 5 minutes; Human Coronavirus, SARS associated Coronavirus and Avian Influenza A with a wet time of 2 minutes; Rhinovirus type 39 and Rotavirus with a wet time of 3 minutes; HIV-1 (associated with AIDS) with a wet time of 1 minute (60 seconds); and Norovirus (Norwalk Virus), Feline Calcivirus and Rabies Virus with a wet time of 30 seconds according to the E.P.A. Pesticide Assessment Guidelines, Subdivision G: Product Performance, Section 91-2 (f), and Section 91-30 (d), (e), November, 1982, modified in the presence of 5% organic serum.

\*\*\* EPA believes, based on available scientific information, that the currently registered influenza A virus products will be effective against the 2009-H1N1 flu strain and other influenza A virus strains on hard, non-porous surfaces.

<http://www.epa.gov/oppad001/influenza-disinfectants.html>